

THE MV FORMALISM FOR IBL_∞ - AND BV_∞ -ALGEBRAS

MARTIN MARKL AND ALEXANDER A. VORONOV

*Однажды чукча принёс в редакцию свой роман. Редактор прочёл и говорит:
 — Понимаете ли, слабовато... Вам бы классику почитать.
 Вы Тургенева читали? А Толстого? А Достоевского?
 — Однако, нет: чукча — не читатель, чукча — писатель.
 Russian folklore*

ABSTRACT. We develop a new formalism for the Quantum Master Equation $\Delta e^{S/\hbar} = 0$ and the category of IBL_∞ -algebras and simplify some homotopical algebra arising in the context of oriented surfaces with boundary. We introduce and study a category of MV-algebras, which, on the one hand, contains such important categories as those of IBL_∞ -algebras and L_∞ -algebras, and on the other hand, is homotopically trivial, in particular allowing for a simple solution of the quantum master equation. We also show that IBL_∞ -morphisms are closed under composition, a nontrivial property, which seems to be taken for granted in the literature.

CONTENTS

1. Introduction	1
2. MV-algebras	4
3. Arithmetic of convolution product	7
4. Category of MV-algebras	11
5. A composition formula and IBL_∞ -algebras	15
6. The Quantum Master Equation	19
References	23

1. INTRODUCTION

Recent developments in String Topology, Symplectic Field Theory, and Lagrangian Floer Theory have led to a new wave of homotopical algebra, heavily burdened by formulas that seem overwhelming to the eye of an unpretentious mathematician, see [6, 5]. The algebra is associated to a homotopy version of involutive Lie bialgebras, or IBL_∞ -algebras. The

Key words and phrases. MV-algebra, IBL_∞ -algebra, Master Equation, Transfer.

The first author was supported by the Eduard Čech Institute P201/12/G028 and RVO: 67985840. The second author was supported by the World Premier International Research Center Initiative (WPI Initiative), MEXT, Japan, and a Collaboration grant from the Simons Foundation (#282349).

algebraic structure in question is governed by surfaces with boundary. It arises from multilinear operators that correspond to diffeomorphism classes of connected, oriented surfaces of various genera and with various numbers of labeled boundary components, separated into “inputs” and “outputs,” like in Topological Quantum Field Theory (TQFT). Since such a diffeomorphism class is determined by the genus and the number of inputs and outputs, the algebraic structure is determined by linear operators: $l_{m,n}^g : S^m(U) \rightarrow S^n(U)$, $g \geq 0$, $m, n \geq 1$, between symmetric powers of a graded vector space U . However, unlike in TQFT, the correspondence is not supposed to be a functor: $l_{1,1}^0 : U \rightarrow U$ has to be a differential on U rather than the identity, and $l_{m,n}^g$ must be a null-homotopy for the sum of all possible gluings of two surfaces along part of their inputs and outputs resulting in a surface of genus g with m inputs and n outputs, see [7, 17, 5]. These operations, $l_{m,n}^g$ may be collected nicely into a generating function $\Delta : S(U)[[\hbar]] \rightarrow S(U)[[\hbar]]$, called a BV (or BV_∞) operator, which is in fact an odd differential operator such that $\Delta^2 = 0$, see [7, 5, 16]. From the operadic perspective, we are talking about the differential graded (dg) dual notion to the notion of a TQFT, namely an algebra over the dg dual properad to the Frobenius algebra properad, which is Koszul dual to the involutive Lie bialgebra properad, see [7, 4]. The standard operadic yoga applies and shows that the algebraic structure is a canonical homotopy version of the structure of an involutive Lie bialgebra, producing an (almost) canonical name: that of an IBL_∞ -algebra.

In concrete geometric contexts, the construction of an IBL_∞ structure involves a number of choices of geometric data, such as an almost complex structure on the target space, a Riemannian metric on the surface, &c., and showing the independence of the algebraic structure on the choices requires the notion of an IBL_∞ -morphism. That notion is quite elaborate algebraically, see [5], but may be packed nicely in a generating function, involving the notion of an “exponential” of a linear map $S(U') \rightarrow S(U'')$ between symmetric algebras. The exponential was defined in [6, 5] by an explicit formula, see (15), bearing a certain resemblance to the exponential power series, as well as respecting the exponentials of elements in these symmetric algebras, see [2], Example 13 and Theorem 44.

The current work came out of the authors’ realization that the exponential of a map was a true exponential in a (graded) commutative algebra given by the convolution product, see (5) and (15). This led to a significant simplification of dealings with the category of IBL_∞ -algebras as well as to a “larger” category of MV-algebras, which we present in this paper. We show that the category of MV-algebras is equivalent to a certain category of pointed complexes. Thus, the category of MV-algebras is rather trivial as a category, but this does not prevent it from having room for such highly nontrivial subcategories as those of L_∞ - and IBL_∞ -algebras. This categorical triviality may be regarded as homotopy triviality, presenting itself through the observation that the quantum master equation (1) in an MV-algebra is just a cocycle condition. We also show that IBL_∞ -morphisms are closed under composition, a nontrivial property, which seems to be taken for granted in the literature.

The theme of the current article may also be viewed as the study of the quantum master equation (QME)

$$(1) \quad \Delta e^{S/\hbar} = 0,$$

which originates in the BV formalism, see, e.g., [13], and takes place in a dg BV- or, more generally, (hereafter, commutative) BV_∞ -algebra. IBL_∞ -algebras provide an important, but not exhaustive class of examples of BV_∞ -algebras: those for which the underlying commutative algebra is free, see Examples 9 and 10. The QME may be viewed as an $\hbar$ -deformation of the Maurer-Cartan equation $dS + \frac{1}{2}[S, S] = 0$, cf. Theorem 38. Thus, the study of solutions of the QME must be related to a quantization of deformation theory, cf. [17], in which the question of homotopy invariance of solutions with respect to weak equivalences would be one of the primary problems. The homotopy category of dg BV- and BV_∞ -algebras should be one of the working notions of quantized deformation theory. Cieliebak and Latschev [6] defined the notion of a BV_∞ -morphism, only when the source was free as a commutative algebra, i.e. derived from an IBL_∞ -algebra. Such a mismatch of sources and targets prevents BV_∞ -algebras from forming a category. MV-morphisms described in this paper include such BV_∞ -morphisms and, on the other hand, form a category. Moreover, the QME makes perfect sense in the more general context of MV-algebras, see Section 6, which makes them arguably a better candidate for providing a background for quantum deformation theory. The main results of the paper in this direction are Theorem 39, which may be interpreted as a representability theorem for the functor of solutions of the QME, see Remark 40, and a description of the transfer of solutions of QME, Theorem 44.

Disclaimer. It may appear that our choice of terminology goes against the good old mathematical tradition of being modestly egocentric. We should assure the reader that, on the contrary, we have chosen to be modest about the mathematical content of the paper and, at the same time, allowed ourselves to be somewhat egocentric in such a minor, cosmetic issue as terminology.

Conventions. The symbol k will denote a fixed field k of characteristic zero and $\otimes$ the tensor product over k . We will denote by id_X or simply by id when X is understood, the identity endomorphism of an object X (vector space, algebra, &c.). We will sometimes denote the product of elements a and b of an algebra using the explicit name of the multiplication (typically $\mu(a, b)$), sometimes, when the meaning is clear from the context, by $a \cdot b$, or simply by ab .

For a graded k -vector space V and a complete local commutative ring R with the residue field k and the maximal ideal $\mathfrak{m}$ we denote by $V \widehat{\otimes} R$ the completed tensor product $\varprojlim_n V \otimes R / V \otimes \mathfrak{m}^n$. In the particular case when R is the formal power series ring $k[[\hbar]]$ in $\hbar$ we abbreviate, as usual, $V \widehat{\otimes} k[[\hbar]]$ by $V[[\hbar]]$. We also abbreviate $k((\hbar)) := k[[\hbar]][\hbar^{-1}]$ and $V((\hbar)) := V \widehat{\otimes} k((\hbar))$. The degree of a homogeneous element $v \in V$ is denoted by $|v|$.

Recall that, for graded indeterminates $u_1, \dots, u_n$ and a permutation $\sigma \in \Sigma_n$, the *Koszul sign* $\epsilon(\sigma) = \epsilon(\sigma; u_1, \dots, u_n) \in \{+1, -1\}$ is defined by

$$u_1 \odot \cdots \odot u_n = \epsilon(\sigma; u_1, \dots, u_n) \cdot u_{\sigma(1)} \odot \cdots \odot u_{\sigma(n)}$$

which has to be satisfied in the free graded commutative algebra $S(u_1, \dots, u_n)$ generated by $u_1, \dots, u_n$. If $k \geq 1$ and $a_1, \dots, a_k$ are non-negative integers such that $a_1 + \cdots + a_k = n$, then an $(a_1, \dots, a_k)$ -*unshuffle* of n elements is a permutation $\sigma \in \Sigma_n$ such that

$$\sigma(1) < \cdots < \sigma(a_1), \sigma(a_1 + 1) < \cdots < \sigma(a_1 + a_2), \dots, \sigma(n - a_k + 1) < \cdots < \sigma(n).$$

The subset of all $(a_1, \dots, a_k)$ -unshuffles will be denoted by $\Sigma_{a_1, \dots, a_k} \subset \Sigma_n$.

Acknowledgment. The authors are indebted to D. Bashkirov for useful remarks on an early draft, and to J. Latschev for sharing a preliminary version of his paper with K. Cieliebak and K. Fukaya. The first author also wishes to express his thanks to M. Doubek, B. Jurčo and I. Sachs for introducing him to the jungle of IBL_∞ -algebras, and acknowledge the hospitality of the University of Minnesota where he held the position of an Distinguished Ordway Visitor during the last stages of the work on this article. The second author gratefully acknowledges support from the Graduate School of Mathematical Sciences, the University of Tokyo, and the Simons Center for Geometry and Physics, Stony Brook University, at which some of the research for this paper was performed.

2. MV-ALGEBRAS

Definition 1. Let R be a complete local Noetherian commutative ring with the residue field k and a maximal ideal $\mathfrak{m}$. An *MV-algebra** over R is a quadruple $V = (V, \mu, \delta, \Delta)$ consisting of

- (i) a unital graded associative commutative k -algebra (V, μ) ,
- (ii) a *conilpotent* counital graded coassociative cocommutative k -coalgebra (V, δ) , and
- (iii) a continuous degree $+1$ R -linear map $\Delta : V \widehat{\otimes} R \rightarrow V \widehat{\otimes} R$ such that $\Delta^2 = 0$ and $\Delta(1) = 0$.

We require that the unit map $\eta : k \rightarrow V$ for (V, μ) is a coaugmentation for (V, δ) , and the counit $\epsilon : V \rightarrow k$ is an augmentation for (V, μ) . In most situations, R will either be the ground field k or a power series ring $k[[\hbar]]$.

Remark 2. Let $I = \text{Ker}(\epsilon)$ be the augmentation ideal of (V, μ) . The *reduced diagonal* $\bar{\delta}$ is defined by the standard formula

$$\bar{\delta}(v) := \delta(v) - (v \otimes 1) - (1 \otimes v)$$

for $v \in I$, while $\bar{\delta}(1) := 0$. We iterate $\bar{\delta}$ by putting $\bar{\delta}^{[0]}(v) := v$ and

$$\bar{\delta}^{[k]}(v) := (\bar{\delta} \otimes id^{\otimes k-1}) \bar{\delta}^{[k-1]}(v)$$

*Abbreviating Markl-Voronov. Not to be mistaken with MV-algebras occurring in many-valued logic.

for $k \geq 1$. The *conilpotency* of (V, δ) means that for each $v \in V$, $\bar{\delta}^{[k]}(v) = 0$ for k large enough. Notice that we *do not* require any compatibility between μ and δ , which would be required of (V, μ, δ) to be a bialgebra, a Frobenius algebra, &c.

Remark 3. Each MV-algebra over k can be considered as an MV-algebra over R after the R -linear extension of Δ . In this manner we obtain precisely those MV-algebras over R for which $\Delta(V) \subset V$.

Example 4. Every augmented unital commutative associative k -algebra A is an MV-algebra over k with the comultiplication defined by $\delta(1) := 1 \otimes 1$, while $\delta(v) := v \otimes 1 + 1 \otimes v$ for v in the augmentation ideal of A , and $\Delta := 0$.

Example 5. Dually, a conilpotent cocommutative coassociative k -coalgebra (V, δ) with a counit $\epsilon : V \rightarrow k$ and a coaugmentation $\eta : k \rightarrow V$ is an MV-algebra over k with a unit η , augmentation ϵ and multiplication given by $1 \cdot v = v \cdot 1 := v$ for each $v \in V$, and $v' \cdot v'' = 0$ for $v', v'' \in \text{Ker}(\epsilon)$. We again put $\Delta := 0$.

Example 6. Each graded k -vector space V equipped with linear maps $\epsilon : V \rightarrow k$ and $\eta : k \rightarrow V$ such that $\epsilon \circ \eta = \text{id}$ bears the ‘supertrivial’ unital algebra and counital coalgebra structures defined similarly as in the above two examples. Such V is therefore a ‘supertrivial’ MV-algebra with any R -linear $\Delta : V \hat{\otimes} R \rightarrow V \hat{\otimes} R$ satisfying $\Delta^2 = 0$ and $\Delta \circ \eta = 0$.

Example 7. Let R be such that the quotient $R/\mathfrak{m}^p$ is a finite-dimensional k -vector space for each $p \geq 1$. The continuous k -linear dual R^* , with the comultiplication given by the dual of the multiplication in R , is conilpotent by the completeness of R . In particular, R^* itself with $\Delta := 0$ and trivial multiplication, as in Example 5, is an MV-algebra over k .

Example 8. A less trivial example of an MV-algebra over k is obtained by taking $V := S(U)$, the (graded) symmetric, or polynomial, algebra generated by a graded vector space U , with the standard bialgebra structure, and Δ a degree 1 coderivation with $\Delta^2 = 0$ and $\Delta(1) = 0$. Such a structure is a disguise of a *strongly homotopy Lie* (L_∞ -) algebra [12, Theorem 2.3].

Let A be a unital associative commutative algebra and $\Delta : A \rightarrow A$ be a k -linear map. For $n \geq 0$, consider the iterated graded commutators

$$(2) \quad [[\dots [\Delta, L_{a_1}], \dots], L_{a_n}],$$

L_a denoting the operator of left multiplication by $a \in A$. By convention, we just set the commutator of Δ with $n = 0$ left-multiplication operators to be Δ . We call an operator Δ an *order $\leq k$ differential operator* if the iterated commutator with any $k + 1$ left-multiplication operators vanishes.

Now suppose that $\Delta(1) = 0$. Define

$$(3) \quad \Phi_n^\Delta(a_1, \dots, a_n) := [[\dots [\Delta, L_{a_1}], \dots], L_{a_n}](1).$$

In particular, $\Phi_0^\Delta = 0$ and $\Phi_1^\Delta = \Delta$. If $\Phi_n^\Delta = 0$ for $n > k$, the operator Δ is called an *order k derivation* [15, Section 1.2]). Notice that first-order derivations are vector fields (derivations).

Example 9. Recall [11, Definition 7] that a (*commutative*) BV_∞ -*algebra* consists of a unital graded commutative associative algebra A and a $k[[\hbar]]$ -linear degree 1 map $\Delta : A[[\hbar]] \rightarrow A[[\hbar]]$ such that $\Delta^2 = 0$ and $\Delta(1) = 0$.[†] One moreover requires that Δ decomposes into a sum

$$(4) \quad \Delta = \Delta_1 + \hbar \Delta_2 + \hbar^2 \Delta_3 + \cdots,$$

where $\Delta_k : A \rightarrow A$ is a k -linear order $\leq k$ differential operator on A . If we assume that A has an augmentation as a graded commutative algebra and equip A with the comultiplication δ constructed in Example 4, then $A = (A, \mu, \delta, \Delta)$ becomes an MV-algebra over $k[[\hbar]]$. The most common degree convention is $|\hbar| = 2$, which is implicit in [11] and explicit in [2], but one can consider $\hbar$ of degree zero, as in [16], or arbitrary even degree, as in [6, 5]. To comply with our convention that the ground ring R is not graded, we set $|\hbar| = 0$.

Example 10. Let $A := S(U)$ be the symmetric algebra generated by a graded vector space U . An IBL_∞ -*algebra* structure on U [5], [16, §4.2] is given by a degree 1, $k[[\hbar]]$ -linear map $\Delta : A[[\hbar]] \rightarrow A[[\hbar]]$ satisfying $\Delta^2 = 0$, $\Delta(1) = 0$, which decomposes as in (4). In other words, an IBL_∞ -algebra is a BV_∞ -algebra whose underlying augmented commutative algebra is $S(U)$. If we equip $S(U)$ with the standard bialgebra structure, $(S(U), \mu, \delta, \Delta)$ will be another example of an MV-algebra over $k[[\hbar]]$.

Remark 11 (Geometric interpretation, see Table 1). One can think of an MV-algebra V over R as the algebra of functions on a family, that is to say a fiber bundle, $X \rightarrow B$ of graded manifolds endowed with a square-zero differential operator Δ of degree one on X over B , thereby, generalizing the notion of a differential graded manifold, when the operator happens to be a derivation linear over functions on B . This fiber bundle has a section $B \rightarrow X$, thought of as a family of basepoints, and the space of distributions on X over B is provided with a graded commutative product, not necessarily related to multiplication of functions on X . The only relationship between the products on functions and distributions we place is that the projection $X \rightarrow B$ and the section $B \rightarrow X$ must provide an augmentation and a unit, respectively, for multiplication of distributions. This geometric object may be called an MV-*manifold*. A typical situation is when the fiber bundle is trivial: $F \times B \rightarrow B$, and one can think of an MV-manifold as a B -parameterized family of differential operators on F . See more on geometric analogies in Table 1.

Since the definition is symmetric, up to technical details, with respect to the algebra-coalgebra structures, we can dually consider the graded algebra structure on the k -linear dual V^* defined by the graded coalgebra structure on V and think of V^* as the algebra of functions on a family $X^* \rightarrow B$ of “formal pointed graded manifolds” endowed with a square-zero, odd “differential operator” over B and the structure of a graded commutative algebra on the space of distributions on X^* , along with similar compatibility conditions between the units and counits. We will call such geometric objects *dual MV-manifolds*. Depending on the situation, either interpretation could be preferable. For instance, in Example 8, the

[†]The continuity of Δ is automatic because $k[[\hbar]]$ is regular, see [14, Proposition 1.11].

Algebra	Geometry
Ring R	Base manifold B
Graded algebra $V \hat{\otimes} R$	Graded manifold X
Unit $\eta : R \rightarrow V \hat{\otimes} R$	Fiber bundle $X \rightarrow B$
Operator Δ	Differential operator on X over B
Augmentation $\epsilon : V \hat{\otimes} R \rightarrow R$	Section $B \rightarrow X$
The R -linear dual $\text{Lin}_R(V \hat{\otimes} R, R)$	Distributions on X over B
Comultiplication on $V \hat{\otimes} R$	Multiplication of distributions on X

TABLE 1. Dictionary of Geometric Analogies

MV-algebra $V = S(U)$ corresponding to an L_∞ -algebra $U[-1]$, where $U[-1]$ denotes an appropriate degree shift, is usually interpreted as a *formal differential graded manifold* U . This is an example of a dual MV-manifold $X = \text{Spec } V^*$. On the other hand, in Example 9, the MV-algebra A may rather be interpreted as a BV_∞ -manifold $\text{Spec } A$, i.e. a family of graded manifolds with an odd, square-zero differential operator over $\text{Spec } k[[\hbar]]$. This is an example of an MV-manifold $\text{Spec } A$.

Geometric objects of this nature may arise in various situations, starting from a graded manifold or graded scheme X provided with extra structure, such as those in the following examples:

- X is a graded “abelian Lie group,” see Examples 8 and 10, in which X is moreover a vector space (or a vector bundle over B).
- X has a “volume density” inducing a graded Frobenius algebra structure on the space of functions on X .
- X has a basepoint x_0 and is provided with a “trivial” multiplication law on distributions so that distributions vanishing on constants multiply to zero and the delta function δ_{x_0} serves as a unit, see Examples 4 and 9. In particular, the delta functions of points multiply as follows: $\delta_x \cdot \delta_y := \delta_x + \delta_y - \delta_{x_0}$ for $x, y \in X$. In this case X^* is a point with a ring V^* of functions such that the maximal ideal I^* of this point has trivial multiplication, $(I^*)^2 = 0$.
- Dually, X may have the “infinitesimal” geometry of a point enriched with a ring V of functions such that the maximal ideal I has trivial multiplication $I^2 = 0$, whereas the space V^* of distributions may have an interesting multiplication, reflected in some nontrivial geometry of X^* , as in Example 5.

3. ARITHMETIC OF CONVOLUTION PRODUCT

Let $V' = (V', \mu', \delta', \Delta')$, $V'' = (V'', \mu'', \delta'', \Delta'')$ be two MV-algebras, and let $\text{Lin}_k(V', V'')$ be the set of degree-0 k -linear maps between their underlying spaces. It is well-known [9, §III.3]

that the comultiplication δ' together with the multiplication μ'' induces on $\text{Lin}_k(V', V'')$ the structure of a unital commutative associative augmented algebra, via the *convolution product* $\star$. Explicitly

$$(5) \quad f \star g := \mu''(f \otimes g)\delta'$$

for $f, g \in \text{Lin}_k(V', V'')$. The unit $\mathbf{e} = \mathbf{e}_{V', V''}$ for $\star$ is the composition $\eta'' \circ \epsilon'$ of the augmentation of V' with the unit of V'' . The algebra $\text{Lin}_k(V', V'')$ is augmented by the map that sends $f \in \text{Lin}_k(V', V'')$ to $\epsilon''(f(1)) \in k$. Notice that

$$(6) \quad \mathbf{e}_{V'', V'''} \circ \mathbf{e}_{V', V''} = \mathbf{e}_{V', V'''}, \quad \mathbf{e}_{V', V''} \circ \eta' = \eta'', \quad \epsilon'' \circ \mathbf{e}_{V', V''} = \epsilon'.$$

The above constructions clearly extend by R -linearity to the space $\text{Lin}_R(V' \hat{\otimes} R, V'' \hat{\otimes} R)$ of continuous R -linear maps which we will, for brevity, denote by $\text{Lin}_R(V', V'')$ believing that the reader will not be too confused by this shorthand.

Let $\mathfrak{m}$ be the maximal ideal of R . Denote by $\text{Lin}_R^0(V', V'')$ the subset of $\text{Lin}_R(V', V'')$ consisting of continuous R -linear maps $f : V' \hat{\otimes} R \rightarrow V'' \hat{\otimes} R$ such that

$$(7) \quad \text{Im}(f \circ \eta') \subset V'' \hat{\otimes} \mathfrak{m}.$$

Notice that for $f, g \in \text{Lin}_R^0(V', V'')$,

$$(8) \quad f \star g \equiv \mu''(f \otimes g)\bar{\delta}' \pmod{V'' \hat{\otimes} \mathfrak{m}},$$

where $\bar{\delta}'$ is the reduced diagonal associated to δ' , and moreover $\text{Lin}_R^0(V', V'')$ is an ideal in $\text{Lin}_R(V', V'')$. We leave to prove as an exercise that the conilpotency of δ' together with the completeness of R implies:

Lemma 12. *All power series in elements of $\text{Lin}_R^0(V', V'')$ converge.*[§]

In particular, for $f \in \text{Lin}_R^0(V', V'')$ it makes sense to take

$$\exp(f) := \mathbf{e} + f + \frac{f^2}{2!} + \frac{f^3}{3!} + \cdots \in \text{Lin}_R(V', V'') \quad (\text{powers w.r. to the } \star\text{-product}).$$

Notice that, while $\exp(f) \notin \text{Lin}_R^0(V', V'')$, clearly $\exp(f) - \mathbf{e} \in \text{Lin}_R^0(V', V'')$.

If $g \in \text{Lin}_R^0(V', V'')$, then $\mathbf{e} + g \in \text{Lin}_R(V', V'')$ and, thanks to (6), we have

$$(\mathbf{e} + g) \circ \eta' \equiv \eta'' \pmod{V'' \hat{\otimes} \mathfrak{m}}$$

and the power series

$$\log(\mathbf{e} + g) := g - \frac{g^2}{2} + \frac{g^3}{3} - \cdots \in \text{Lin}_R^0(V', V'')$$

converges. For $f, g \in \text{Lin}_R^0(V', V'')$, one clearly has the expected equalities

$$\log(\exp(f)) = f, \quad \exp(\log(\mathbf{e} + g)) = \mathbf{e} + g.$$

[‡]We used the same symbol for the map η' and its R -linear extensions. We keep this convention throughout the paper.

[§]Convergence is always understood as convergence in the $\mathfrak{m}$ -adic topology.

Example 13. Let $G(V') := \{v \in V' \mid \delta'(v) = v \otimes v\}$ be the subset of group-like elements in V' , and $v \in G(V') \widehat{\otimes} R$. Then, for any $f, g \in \text{Lin}_R(V', V'')$,

$$(f \star g)(v) = \mu''(f \otimes_R g) \delta'(v) = \mu''(f \otimes_R g)(v \otimes_R v) = \mu''(f(v), g(v)),$$

so $(f \star g)(v)$ is the ‘actual’ product of the elements $f(v)$ and $g(v)$ in the algebra V'' . Consequently, for $f \in \text{Lin}_R^0(V', V'')$ and $v \in G(V') \widehat{\otimes} R$,

$$\exp(f)(v) = e^{f(v)},$$

the ‘actual’ exponential of $f(v)$ in the algebra V'' ; similarly for the logarithm.

Proposition 14. *Given two MV-algebras V' and V'' and $f \in \text{Lin}_R^0(V', V'')$, suppose that V' is in fact a bialgebra, i.e. δ' is an algebra morphism. Then, for any $v_1, v_2 \in V'$,*

$$(9) \quad \exp(f)(v_1 v_2) \equiv \exp(f)(v_1) \exp(f)(v_2) \pmod{(V'' \widehat{\otimes} \mathbf{m}, f(I'^2))},$$

where I' is the augmentation ideal of V' . In other words, $\exp(f)$ is an algebra morphism modulo the ideal in $V'' \widehat{\otimes} R$ generated by $V'' \widehat{\otimes} \mathbf{m}$ and $f(I'^2)$.

Proof. Since (9) obviously holds if v_1 or v_2 are proportional to $1 \in V'$, we will assume that $v_1, v_2 \in I'$. In this case, since $\epsilon'(v_1) = \epsilon'(v_2) = 0$, $\exp(f)(v_1) \exp(f)(v_2)$ is the product

$$(10) \quad \left(f(v_1) + \frac{1}{2}f^2(v_1) + \frac{1}{3}f^3(v_1) + \cdots\right) \left(f(v_2) + \frac{1}{2}f^2(v_2) + \frac{1}{3}f^3(v_2) + \cdots\right)$$

while

$$(11) \quad \exp(f)(v_1 v_2) = f(v_1 v_2) + \frac{1}{2}f^2(v_1 v_2) + \frac{1}{3}f^3(v_1 v_2) + \cdots.$$

As δ' is, by assumption, an algebra morphism, one has

$$\begin{aligned} \delta'(v_1 v_2) &= \delta'(v_1) \delta'(v_2) = (v_1 \otimes 1 + \bar{\delta}'(v_1) + 1 \otimes v_1)(v_2 \otimes 1 + \bar{\delta}'(v_2) + 1 \otimes v_2) \\ &= v_1 \otimes v_2 + (-1)^{|v_1| \cdot |v_2|} v_2 \otimes v_1 + v_1 v_2 \otimes 1 + 1 \otimes v_1 v_2 + \bar{\delta}'(v_1) \bar{\delta}'(v_2) \\ &\quad + (1 \otimes v_1) \bar{\delta}'(v_2) + (v_1 \otimes 1) \bar{\delta}'(v_2) + \bar{\delta}'(v_1)(1 \otimes v_2) + \bar{\delta}'(v_1)(v_2 \otimes 1). \end{aligned}$$

Notice that $\mu'' \circ (f \otimes f)$ applied to the terms on the right-hand side of the above equation vanishes modulo $(V'' \widehat{\otimes} \mathbf{m}, f(I'^2))$ everywhere except at the terms $v_1 \otimes v_2$ and $v_2 \otimes v_1$, thus

$$f^2(v_1 v_2) = \mu''(f \otimes f) \delta'(v_1 v_2) \equiv \mu''(f(v_1) \otimes f(v_2) + (-1)^{|v_1| \cdot |v_2|} f(v_2) \otimes f(v_1)) = 2f(v_1) f(v_2)$$

modulo $(V'' \widehat{\otimes} \mathbf{m}, f(I'^2))$. Similarly we obtain that

$$f^3(v_1 v_2) = \mu''(f^2 \otimes f) \delta'(v_1 v_2) \equiv 3f^2(v_1) f(v_2) + 3f(v_1) f^2(v_2) \pmod{(V'' \widehat{\otimes} \mathbf{m}, f(I'^2))}$$

and, inductively,

$$f^n(v_1 v_2) \equiv \sum_{1 \leq i \leq n-1} \binom{n}{i} f^i(v_1) f^{n-i}(v_2) \pmod{(V'' \widehat{\otimes} \mathbf{m}, f(I'^2))}.$$

This formula makes the verification that the product (10) equals (11) modulo $(V'' \widehat{\otimes} \mathbf{m}, f(I'^2))$ obvious. $\square$

Corollary 15. *Under the assumptions of Proposition 14, suppose also that $f(I'^2) \subset V'' \widehat{\otimes} \mathfrak{m}$. Then, for any $v_1, v_2 \in V'$,*

$$(12) \quad \exp(f)(v_1 v_2) \equiv \exp(f)(v_1) \exp(f)(v_2) \pmod{V'' \widehat{\otimes} \mathfrak{m}}.$$

Thus, $\exp(f)$ is an algebra morphism modulo $V'' \widehat{\otimes} \mathfrak{m}$.

Example 16. Let us show that Proposition 14 cannot be strengthened. Take $V' = V'' = \mathfrak{k}$ with the obvious bialgebra structure. Then $f \in \text{Lin}_R^0(\mathfrak{k}, \mathfrak{k})$ is determined by $\alpha := f(1) \in \mathfrak{m}$. By definition, $\exp(f)(v) = ve^\alpha$ for $v \in \mathfrak{k}$, so we have

$$\exp(f)(v_1 v_2) = v_1 v_2 e^\alpha$$

while

$$\exp(f)(v_1) \exp(f)(v_2) = v_1 v_2 e^{2\alpha}.$$

The map $\exp(f) : \mathfrak{k} \rightarrow \mathfrak{k}$ is in this case an algebra morphism only modulo the ideal generated by $\alpha \in \mathfrak{m}$.

As the second example, take $V' = V'' = S(U)$ with the standard bialgebra structure, $R := \mathfrak{k}$, and assume that $f : S(U) \rightarrow S(U)$ is such that $f|_{S^n(U)} \neq 0$ only for $n = 2$. One then easily calculates that, for $v_1, v_2 \in U$,

$$\exp(f)(v_1) = \exp(f)(v_2) = 0$$

while

$$\exp(f)(v_1 v_2) = f(v_1 v_2).$$

The map $\exp(f) : S(U) \rightarrow S(U)$ is an algebra morphism only modulo $f(S^{\geq 2}(U))$.

Assume now that $f|_{S^n(U)} \neq 0$ only for $n = 1$. Then $f(S^{\geq 2}(U)) = 0$ and, since $R = \mathfrak{k}$, also $\mathfrak{m} = 0$. Formula (9) therefore asserts that $\exp(f)$ is an algebra morphism. We leave as an exercise to verify that indeed $\exp(f)$ provides the unique extension of a linear map $U \rightarrow S(U)$ into an algebra morphism.

Remark 17. Proposition 14 and Corollary 15 provide sufficient conditions for $\exp(f)$ to be a perturbation of an algebra morphism. See also Examples 25 and 26, in which $\exp(f)$ will be an $\hbar$ -perturbation of an algebra morphism. In the general case, one can view $\exp(f)$ as a generalized algebra morphism.

Example 18. Let us describe the exponential of an R -linear map $f \in \text{Lin}_R^0(S(U), B)$, where $S(U) = (S(U), \delta, \eta)$ is the symmetric algebra generated by a graded vector space U with the standard cocommutative coassociative counital comultiplication, and $B = (B, \mu, 1)$ a commutative associative unital algebra.

Denoting, as usual, by $\delta^{[k-1]} : S(U) \rightarrow S(U)^{\otimes k}$ the diagonal iterated $(k-1)$ -times and, likewise, by $\mu^{[k-1]} : B^{\otimes k} \rightarrow B$ the R -linear extension of the iterated multiplication in B , we have, for $k \geq 1$ and f as above

$$(13) \quad f^k = \mu^{[k-1]} \circ f^{\otimes k} \circ \delta^{[k-1]}.$$

It is easy to verify that the $(k-1)$ -times iterated diagonal on the product $u_1 \odot \cdots \odot u_n \in S^n(U)$ of $u_1, \dots, u_n \in U$ equals

$$(14) \quad \delta^{[k-1]}(u_1 \odot \cdots \odot u_n) = \sum \frac{\epsilon(\sigma)}{a_1! \cdots a_k!} [u_{\sigma(1)} \odot \cdots \odot u_{\sigma(a_1)}] \otimes \cdots \otimes [u_{\sigma(n-a_k+1)} \odot \cdots \odot u_{\sigma(n)}],$$

where the summation runs over all permutations $\sigma \in \Sigma_k$ and integers $a_1, \dots, a_k \geq 0$ such that $a_1 + \cdots + a_k = n$, and $\epsilon(\sigma)$ is the Koszul sign. Here we also use the convention that if $a_i = 0$, then

$$[u_{\sigma(a_1+\cdots+a_{i-1})} \odot \cdots \odot u_{\sigma(a_1+\cdots+a_i)}] = 1.$$

Evaluating $\delta^{[k-1]}$ in (13) using (14) gives, for $u_1 \odot \cdots \odot u_n \in S^n(U)$ and $n \geq 1$,

$$(15) \quad \exp(f)(u_1 \odot \cdots \odot u_n) = \sum_{k=1}^{\infty} \frac{1}{k!} \sum \frac{\epsilon(\sigma)}{a_1! \cdots a_k!} f(u_{\sigma(1)} \odot \cdots \odot u_{\sigma(a_1)}) \cdots f(u_{\sigma(n-a_k+1)} \odot \cdots \odot u_{\sigma(n)}).$$

We recognize a formula in [6, Section 5]. Notice that, thanks to the commutativity of the multiplication $\odot$ in $S(U)$, this formula can be rewritten as

$$(16) \quad \exp(f)(u_1 \odot \cdots \odot u_n) = \sum_{k=1}^{\infty} \frac{1}{k!} \sum \epsilon(\sigma) f(u_{\sigma(1)} \odot \cdots \odot u_{\sigma(a_1)}) \cdots f(u_{\sigma(n-a_k+1)} \odot \cdots \odot u_{\sigma(n)})$$

where σ runs now over all $(a_1, \dots, a_k)$ -unshuffles of n elements. The calculation is completed by

$$\exp(f)(1) = e^{f(1)},$$

the ‘actual’ exponential of $f(1)$ in $B \hat{\otimes} \mathfrak{m}$.

4. CATEGORY OF MV-ALGEBRAS

In this section we define the category of MV-algebras over R . Firstly, we introduce morphisms:

Definition 19. The space of *(MV-)morphisms* between MV-algebras $V' = (V', \mu', \delta', \Delta')$ and $V'' = (V'', \mu'', \delta'', \Delta'')$ is given by

$$\mathbf{MV}_R(V', V'') := \{f \in \text{Lin}_R^0(V', V'') \mid \Delta'' \circ \exp(f) = \exp(f) \circ \Delta'\}.$$

The categorical composition of $f \in \mathbf{MV}_R(V'', V''')$ with $g \in \mathbf{MV}_R(V', V'')$ is defined as

$$f \diamond g := \log(\exp(f) \circ \exp(g)).$$

The unit endomorphism of an MV-algebra V over R is $1_V := \log(\text{id}_{V \hat{\otimes} R})$.

Theorem 20. $\mathbf{MV}_R$ with the above notion of morphisms forms a category.

Proof. Let us verify the associativity of the categorical composition. By definition,

$$(f \diamond g) \diamond h = \log(\exp(f \diamond g) \circ \exp h) = \log(\exp(f) \circ \exp(g) \circ \exp(h))$$

while

$$f \diamond (g \diamond h) = \log(\exp f \circ \exp(g \diamond h)) = \log(\exp(f) \circ \exp(g) \circ \exp(h)),$$

so $(f \diamond g) \diamond h = f \diamond (g \diamond h)$ as required. To verify the axiom for the categorical identity is also easy; one has

$$f \diamond 1_V = \log(\exp(f) \circ id_{V \widehat{\otimes} R}) = \log(\exp(f)) = f.$$

The identity $1_V \diamond f = f$ is verified in a similar fashion. The last thing which remains to be verified is that $\Delta''' \circ \exp(f \diamond g) = \exp(f \diamond g) \circ \Delta'$ which follows from

$$\Delta''' \circ \exp(f \diamond g) = \Delta''' \circ \exp(f) \circ \exp(g) = \exp(f) \circ \exp(g) \circ \Delta' = \exp(f \diamond g) \circ \Delta'$$

□

Example 21. Let us show that, as stated in [2], the unit endomorphism

$$1_{S(U)} \in Lin_k^0(S(U), S(U))$$

of the IBL_∞ -algebra $S(U)$ recalled in Example 10 is the projection $\pi_1 : S(U) \rightarrow U$ to the space of algebra generators. Since π_1 is the projection to U , it follows from formula (14) for the iterated diagonal that $\pi_1^{\otimes k} \circ \delta^{[k-1]}(u_1 \odot \cdots \odot u_n) \neq 0$ only when $k = n$, in which case (14) gives

$$\mu^{[k-1]} \circ \pi_1^{\otimes k} \circ \delta^{[k-1]}(u_1 \odot \cdots \odot u_n) = \begin{cases} n! (u_1 \odot \cdots \odot u_n) & \text{if } n = k \text{ and} \\ 0 & \text{otherwise.} \end{cases}$$

This readily implies that $\exp(\pi_1) = id_{S(U)}$ i.e. that $1_{S(U)} = \pi_1 = \log(id_{S(U)})$ as claimed.

We recommend as an exercise to verify that $\pi_1 = \log(id_{S(U)})$ directly. It turns out that this equation leads to an interesting combinatorial formula for the unshuffles.

Remark 22 (Geometric interpretation). We can interpret a morphism in $MV_R(V', V'')$ geometrically as a generalized, as in Remark 17, morphism $X'' \rightarrow X'$ of MV-manifolds. Dually, we can think of it as a morphism $X'^* \rightarrow X''^*$ of dual MV-manifolds.

Example 23. Notice that if the reduced diagonal in V' is trivial,

$$\exp(f) \equiv e + f \pmod{V'' \widehat{\otimes} \mathfrak{m}}, \text{ and } \log(e + g) = g \pmod{V'' \widehat{\otimes} \mathfrak{m}}.$$

The category MV_k has a full subcategory consisting of MV-algebras over k with trivial reduced diagonals. The composition rule in this subcategory is given by

$$\begin{aligned} f \diamond g &= \log(\exp(f) \circ \exp(g)) = \log((e + f) \circ (e + g)) \\ &= \log(e \circ e + e \circ g + f \circ e + f \circ g) = f \circ g + e \circ g + f \circ e. \end{aligned}$$

We used the fact that $e \circ e = e$ by (6). It is an instructive exercise to verify that the categorical unit endomorphism is $id - e$. Notice that the ‘expected’ unit id is not even an element of $Lin_R^0(V, V)$.

Restricting to an even smaller subcategory whose morphisms $f \in \text{MV}_R(V', V'')$ satisfy the stronger condition

$$f \circ \eta' = 0, \quad \epsilon'' \circ f = 0,$$

the composition rule $f \diamond g$ becomes the standard composition of morphisms.

Example 24. The category MV_k contains the subcategory L_∞ whose objects are L_∞ -algebras recalled in Example 8 and morphisms are maps $f : S(U') \rightarrow S(U'')$ such that

$$f(1) = 0, \quad \Delta'' \circ \exp(f) = \exp(f) \circ \Delta', \quad \text{and} \quad \text{Im}(f) \subset U''.$$

Such a map automatically belongs to $\text{Lin}_k^0(S(U'), S(U''))$. We leave as an interesting exercise to prove that

$$\exp(f) : S(U') \rightarrow S(U'')$$

is the unique coextension of f into a morphism of counital coalgebras. We conclude that L_∞ is isomorphic to the category of L_∞ -algebras and their (weak) L_∞ -morphisms [12, Remark 5.3].

Example 25. Let us consider IBL_∞ -algebras recalled in Example 10 with $k[[\hbar]]$ -linear maps

$$f : S(U')[[\hbar]] \rightarrow S(U'')[[\hbar]]$$

of the form

$$(17) \quad f = f^{(1)} + \hbar f^{(2)} + \hbar^2 f^{(3)} + \dots$$

such that

$$(18) \quad \begin{aligned} f^{(1)}(1) &= 0, \quad \Delta'' \circ \exp(f) = \exp(f) \circ \Delta', \quad \text{and} \\ \bigoplus_{n > k} S^n(U') &\subset \text{Ker}(f^{(k)}). \end{aligned}$$

In Corollary 33 below we prove that the above structure forms a subcategory IBL_∞ of the category $\text{MV}_{k[[\hbar]]}$ of MV-algebras over $k[[\hbar]]$, c.f. also [6, §5] and [5, Definition 2.8]. One can consider a version of this subcategory with the condition (18) replaced with a “dual” condition:

$$\text{Im}(f^{(k)}) \subset \bigoplus_{1 \leq n \leq k} S^n(U'').$$

This modified subcategory of $\text{MV}_{k[[\hbar]]}$ may be called the category of IBL_∞ -algebras in the sense of [16, §4.3].

Example 26 (Cieliebak-Latschev [6]). A BV_∞ -morphism from a BV_∞ -algebra $(S(U), \Delta')$ of Example 10 to a BV_∞ -algebra (A, Δ'') of Example 9 is an MV-morphism given by a $k[[\hbar]]$ -linear map

$$f : S(U)[[\hbar]] \rightarrow A[[\hbar]]$$

of the form

$$f = f^{(1)} + \hbar f^{(2)} + \hbar^2 f^{(3)} + \dots,$$

such that

$$f^{(1)}(1) = 0, \quad \Delta'' \circ \exp(f) = \exp(f) \circ \Delta', \quad \text{and} \\ \bigoplus_{n>k} S^n(U) \subset \text{Ker}(f^{(k)}).$$

This is a generalization of the notion of an IBL_∞ -morphism of the type (18).

We are going to define a product $V' \otimes V''$ of two MV-algebras $V' = (V', \mu', \delta', \Delta')$ and $V'' = (V'', \mu'', \delta'', \Delta'')$ over R as follows. Its underlying graded vector space is $V' \otimes V''$ and the structure operator is given as $\Delta' \otimes_R \text{id} + \text{id} \otimes_R \Delta''$. The multiplication is defined in the standard way:

$$(v'_1 \otimes v''_1) \cdot (v'_2 \otimes v''_2) := (-1)^{|v''_1||v'_2|} v'_1 \cdot v'_2 \otimes v''_1 \cdot v''_2, \quad v'_1, v'_2 \in V', \quad v''_1, v''_2 \in V'',$$

with the unit given by the map $\eta' \otimes \eta'' : \mathbf{k} \cong \mathbf{k} \otimes \mathbf{k} \rightarrow V' \otimes V''$. The comultiplication is defined as

$$\delta(v' \otimes v'') := \tau_{23}(\delta'(v') \otimes \delta''(v'')), \quad v' \in V', \quad v'' \in V'',$$

where τ_{23} permutes the second and the third factors with the Koszul sign and, finally, $\epsilon' \otimes \epsilon'' : V' \otimes V'' \rightarrow \mathbf{k} \otimes \mathbf{k} \cong \mathbf{k}$ is the counit. The $\otimes$ -product of morphisms $f \in \text{MV}_R(V'_1, V'_2)$ and $g \in \text{MV}_R(V''_1, V''_2)$ is given by the formula

$$f \otimes g := \log(\exp(f) \otimes_R \exp(g)).$$

Proposition 27. *The $\otimes$ -product equips MV_R with a symmetric monoidal structure.*

Proof. Direct verification. □

Proposition 28. *The category MV_R of MV-algebras over R is isomorphic to the category $\widetilde{\text{MV}}_R$ with the same objects and morphisms*

$$\widetilde{\text{MV}}_R(V', V'') := \{\varphi \in \text{Lin}_R(V', V'') \mid \varphi \circ \eta' \equiv \eta'' \text{ mod } V'' \widehat{\otimes} \mathbf{m} \text{ and } \Delta'' \circ \varphi = \varphi \circ \Delta'\}.$$

The categorical composition is the usual composition of maps, and the unit $1_V \in \widetilde{\text{MV}}_R(V, V)$ is the identity $\text{id} : V \rightarrow V$.

Proof. The isomorphism between MV_R and $\widetilde{\text{MV}}_R$ is given by the mutually inverse functors identical on objects and taking a morphism $f \in \text{MV}_R(V', V'')$ to $\exp(f) \in \widetilde{\text{MV}}_R(V', V'')$ and $\varphi \in \widetilde{\text{MV}}_R(V', V'')$ to $\log(\varphi) \in \text{MV}_R(V', V'')$. Let us check that this isomorphism is well-defined.

Notice first that for $\varphi \in \text{Lin}_R(V', V'')$, the condition $\varphi \circ \eta' \equiv \eta'' \text{ mod } V'' \widehat{\otimes} \mathbf{m}$ is equivalent to the condition $(\varphi - \mathbf{e}) \circ \eta' \equiv 0 \text{ mod } V'' \widehat{\otimes} \mathbf{m}$. To see it, recall that $\mathbf{e} = \eta'' \circ \epsilon'$, thus

$$(\varphi - \mathbf{e}) \circ \eta' = \varphi \circ \eta' - \eta'' \circ \epsilon' \circ \eta' = \varphi \circ \eta' - \eta''.$$

Likewise, for $f \in \text{Lin}_R(V', V'')$, $f \circ \eta' \equiv 0 \text{ mod } V'' \widehat{\otimes} \mathbf{m}$ is equivalent to $(f + \mathbf{e}) \circ \eta' \equiv \eta'' \text{ mod } V'' \widehat{\otimes} \mathbf{m}$.

It is now easy to verify, using the definitions of $\exp$ and $\log$, that $\exp(f)$ indeed belongs to $\widetilde{\text{MV}}_R(V', V'')$ and $\log(\varphi)$ to $\text{MV}_R(V', V'')$. The fact that the above correspondence converts the $\diamond$ -composition to the usual one is clear. □

Corollary 29. *The category MV_R of MV-algebras over R is equivalent to the category Chn_R° of pointed complexes over R whose objects are graded vector spaces V with a continuous degree $+1$ R -linear differential $\Delta : V \hat{\otimes} R \rightarrow V \hat{\otimes} R$ and a $\mathbb{k}$ -linear morphism $\eta : \mathbb{k} \rightarrow V$ such that $\Delta \circ \eta = 0$. A morphism between V' and V'' is a chain map $\varphi \in \text{Lin}_R(V', V'')$ such that $\varphi \circ \eta' \equiv \eta'' \pmod{V'' \hat{\otimes} \mathfrak{m}}$.*

Proof. By Proposition 28, it is enough to establish an equivalence between the categories $\widetilde{\text{MV}}_R$ and Chn_R° . Let us construct mutual weak inverses $\square : \widetilde{\text{MV}}_R \rightarrow \text{Chn}_R^\circ$ and $F : \text{Chn}_R^\circ \rightarrow \widetilde{\text{MV}}_R$.

On objects, the functor $\square$ forgets everything except the structure operator Δ and the unit map η . For $(V, \Delta, \eta) \in \text{Chn}_R^\circ$ choose a right inverse ϵ of η and define $F(V, \Delta, \eta)$ the supertrivial MV-algebra as in Example 6. Notice that, for $V', V'' \in \widetilde{\text{MV}}_R$,

$$\text{Chn}_R^\circ(\square V', \square V'') = \widetilde{\text{MV}}_R(V', V'')$$

and, likewise, for $V', V'' \in \text{Chn}_R^\circ$,

$$\widetilde{\text{MV}}_R(FV', FV'') = \text{Chn}_R^\circ(V', V'').$$

We define $\square$ and F to be the identities on morphisms. It is simple to verify that we have constructed mutual weak inverses. $\square$

Remark 30. The definition of the category of MV-algebras can be modified. For instance, we may leave the conilpotency of δ out, but instead of (7) require that $\text{Im}(f) \in V'' \hat{\otimes} \mathfrak{m}$. Likewise, we need not require R to be complete, but then (7) must be replaced by $f \circ \eta' = 0$. In both cases the above constructions remain valid. We may also require $\epsilon \circ \Delta = 0$, drop the condition $\Delta(1) = 0$, or require both conditions simultaneously.

5. A COMPOSITION FORMULA AND IBL_∞ -ALGEBRAS

Let us consider morphisms $g \in \text{Lin}_R^0(S(U'), S(U''))$ and $f \in \text{Lin}_R^0(S(U''), S(U'''))$, where $S(U'), S(U'')$ and $S(U''')$ are symmetric algebras with the standard coalgebra structures. The aim of this section is to give an explicit formula for

$$(19) \quad f \diamond g = \log(\exp(f) \circ \exp(g)) \in \text{Lin}_R^0(S(U'), S(U''')).$$

Further, using this formula, we prove that IBL_∞ -algebras with morphisms (17) form a subcategory of $\text{MV}_{\mathbb{k}[[h]]}$.

Let us formulate some preparatory observations. Each R -linear map

$$h : S(V') \hat{\otimes} R \rightarrow S(V'') \hat{\otimes} R$$

determines a family

$$h_n^m : S^n(V') \rightarrow S^m(V'') \hat{\otimes} R, \quad m, n \geq 0,$$

with h_m^n the composition

$$S^n(V') \hookrightarrow S(V') \xrightarrow{h|_{S(V')}} S(V'') \hat{\otimes} R \twoheadrightarrow S^n(V'') \hat{\otimes} R, \P$$

$\P$ We use the convention that $S^0(V') = S^0(V'') = \mathbb{k}$.

where $\hookrightarrow$ resp. $\twoheadrightarrow$ is the canonical inclusion resp. the canonical projection. Vice versa, each family $\{h_n^m\}_{m,n \geq 0}$ as above such that the sum

$$h|_{S(V')}(x) := \sum_{m \geq 0} h_n^m(x)$$

converges in $S(V'') \widehat{\otimes} R$ for each fixed $n \geq 0$ and $x \in S^n(V')$, determines an R -linear map $h : S(V') \widehat{\otimes} R \rightarrow S(V'') \widehat{\otimes} R$. We will call h_n^m the $\binom{m}{n}$ -component of h . We will describe $f \diamond g$ in terms of its $\binom{m}{n}$ -components.

For natural numbers k, l and non-negative integers $r, s_1, \dots, s_l, j_1, \dots, j_k$ such that

$$j_1 + \dots + j_k = s_1 + \dots + s_l = r$$

we define

$$(20) \quad \Psi_{j_1, \dots, j_k}^{s_1, \dots, s_l} : S^{j_1}(U'') \otimes \dots \otimes S^{j_k}(U'') \longrightarrow S^{s_1}(U'') \otimes \dots \otimes S^{s_l}(U'')$$

to be the k -linear map that sends

$$(u_1'' \odot \dots \odot u_{j_1}'') \otimes \dots \otimes (u_{r-j_k+1}'' \odot \dots \odot u_r'') \in S^{j_1}(U'') \otimes \dots \otimes S^{j_k}(U'')$$

into the sum

$$(21) \quad \sum_{\kappa \in \Sigma_{s_1, \dots, s_l}} \epsilon(\kappa) (u_{\kappa(1)}'' \odot \dots \odot u_{\kappa(s_1)}'') \otimes \dots \otimes (u_{\kappa(r-s_l+1)}'' \odot \dots \odot u_{\kappa(r)}'') \in S^{s_1}(U'') \otimes \dots \otimes S^{s_l}(U'')$$

over the set $\Sigma_{s_1, \dots, s_l}$ of all $(s_1, \dots, s_l)$ -unshuffles κ of $r = s_1 + \dots + s_l$ elements. It is easy to check that (21) is well-defined, i.e. that (21) is invariant under permutations of generators inside the groups

$$(22) \quad \{u_1'', \dots, u_{j_1}''\}, \dots, \{u_{r-j_k+1}'', \dots, u_r''\}.$$

We associate to each $\kappa \in \Sigma_{s_1, \dots, s_l}$ in (21) a graph Γ with two types of vertices, the upper and lower ones. The upper ones are labelled by $1, \dots, l$, the lower ones by $1, \dots, k$. The upper vertex labelled by $b \in \{1, \dots, l\}$ is connected to the lower vertex labeled by $a \in \{1, \dots, k\}$ if and only if κ maps some element of the b th segment of the decomposition

$$\{1, \dots, r\} = \{1, \dots, s_1\} \cup \dots \cup \{r - s_l + 1, \dots, r\}$$

to some element of the a th segment of the decomposition

$$\{1, \dots, r\} = \{1, \dots, j_1\} \cup \dots \cup \{r - j_k + 1, \dots, r\}.$$

Let $c(\kappa)$ be the number of connected components of Γ . We will call $c(\kappa)$ the *connectivity* of κ and say that κ is *connected* if $c(\kappa) = 1$. Denote by ${}^c\Sigma_{s_1, \dots, s_l}^{j_1, \dots, j_k}$ the subset of all $\kappa \in \Sigma_{s_1, \dots, s_l}$ of connectivity c and let finally

$${}^c\Psi_{j_1, \dots, j_k}^{s_1, \dots, s_l} : S^{j_1}(U'') \otimes \dots \otimes S^{j_k}(U'') \rightarrow S^{s_1}(U'') \otimes \dots \otimes S^{s_l}(U'')$$

be the map defined as $\Psi_{j_1, \dots, j_k}^{s_1, \dots, s_l}$ but with the sum (21) restricted to connected $\kappa \in {}^c\Sigma_{s_1, \dots, s_l}^{j_1, \dots, j_k}$.

Proposition 31. *The maps ${}^c\Psi_{j_1, \dots, j_k}^{s_1, \dots, s_l}$, $c \geq 1$, are well-defined and*

$$(23) \quad \Psi_{j_1, \dots, j_k}^{s_1, \dots, s_l} = {}^1\Psi_{j_1, \dots, j_k}^{s_1, \dots, s_l} + {}^2\Psi_{j_1, \dots, j_k}^{s_1, \dots, s_l} + {}^3\Psi_{j_1, \dots, j_k}^{s_1, \dots, s_l} + \dots$$

Proof. Since composing with permutations inside the groups (22) does not change the connectivity of κ , each individual ${}^c\Psi_{j_1, \dots, j_k}^{s_1, \dots, s_l}$ is well-defined. Formula (23) is obvious. $\square$

In what follows we use the same symbols both for the maps (23) and for their R -linear extensions. Therefore, for instance, ${}^1\Psi_{j_1, \dots, j_k}^{s_1, \dots, s_l}$ will also denote an R -linear map

$$(24) \quad {}^1\Psi_{j_1, \dots, j_k}^{s_1, \dots, s_l} : [S^{j_1}(U'') \otimes \dots \otimes S^{j_k}(U'')] \widehat{\otimes} R \rightarrow [S^{s_1}(U'') \otimes \dots \otimes S^{s_l}(U'')] \widehat{\otimes} R.$$

Let $g_n^m : S^n(U') \rightarrow S^m(U'') \widehat{\otimes} R$ be, for $m, n \geq 0$, the $\binom{m}{n}$ -components of g and let

$$f_n^m : S^n(U'') \widehat{\otimes} R \rightarrow S^m(U''') \widehat{\otimes} R$$

be the R -linear extensions of the $\binom{m}{n}$ -components of f . For given natural numbers k, l and non-negative integers $i_1, \dots, i_k, t_1, \dots, t_l$ we define an auxiliary map

$$(f \diamond g)_{i_1, \dots, i_k}^{t_1, \dots, t_l} : S^{j_1}(U'') \otimes \dots \otimes S^{j_k}(U'') \longrightarrow [S^{s_1}(U'') \otimes \dots \otimes S^{s_l}(U'')] \widehat{\otimes} R$$

as the sum

$$\sum \frac{1}{k! \, l!} (f_{s_1}^{t_1} \otimes \dots \otimes f_{s_l}^{t_l}) \circ {}^1\Psi_{j_1, \dots, j_k}^{s_1, \dots, s_l} \circ (g_{i_1}^{j_1} \otimes \dots \otimes g_{i_k}^{j_k})$$

over all non-negative integers $s_1, \dots, s_l, j_1, \dots, j_k$ such that $s_1 + \dots + s_l = j_1 + \dots + j_k$, where ${}^1\Psi_{j_1, \dots, j_k}^{s_1, \dots, s_l}$ is the R -linear extension (24) of the connected component of the map $\Psi_{j_1, \dots, j_k}^{s_1, \dots, s_l}$ in (20).

In the proof of Theorem 32 below we write (25) in the form

$$(25) \quad \sum \frac{1}{k! \, l!} \left(\begin{array}{c} f_{s_1}^{t_1} \otimes \dots \otimes f_{s_l}^{t_l} \\ {}^1\Psi_{j_1, \dots, j_k}^{s_1, \dots, s_l} \\ g_{i_1}^{j_1} \otimes \dots \otimes g_{i_k}^{j_k} \end{array} \right)$$

which would enable us to squeeze formula (28) into a line of finite length. Finally, for $m, n > 0$ define a k -linear map

$$(f \diamond g)_n^m : S^n(U') \rightarrow S^m(U''') \widehat{\otimes} R$$

by the formula

$$(26) \quad (f \diamond g)_n^m(u'_1 \odot \dots \odot u'_n) := \sum \mu^{[l-1]} \epsilon(\sigma) (f \diamond g)_{i_1, \dots, i_k}^{t_1, \dots, t_l} (u'_{\sigma(1)} \odot \dots \odot u'_{\sigma(i_1)}) \otimes \dots \otimes (u'_{\sigma(n-i_k+1)} \odot \dots \odot u'_{\sigma(n)}),$$

where the summation runs over all natural numbers k, l , non-negative integers $i_1, \dots, i_k, t_1, \dots, t_l$ such that

$$i_1 + \dots + i_k = n \quad \text{and} \quad t_1 + \dots + t_l = m,$$

and over all $(i_1, \dots, i_k)$ -unshuffles $\sigma \in \Sigma_{i_1, \dots, i_k}$. In (26),

$$\mu^{[l-1]} : [S^{t_1}(U''') \otimes \dots \otimes S^{t_l}(U''')] \widehat{\otimes} R \longrightarrow S^m(U''') \widehat{\otimes} R$$

is the R -linear extension of the multiplication in $S(U''')$ iterated $(l-1)$ -times.

Theorem 32. *The $\binom{m}{n}$ -part of the composition $f \diamond g \in \text{Lin}_R^0(S(U'), S(U'''))$ is given by formula (26).*

Proof. Our strategy will be to show that the exponential of the map whose $\binom{m}{n}$ -parts are given by (26) equals $\exp(f) \circ \exp(g)$. Using (16) we find the following expression for the $\binom{m}{n}$ -components of the composition $\exp(f) \circ \exp(g)$:

$$\begin{aligned} & (\exp(f) \circ \exp(g))_n^m (u'_1 \odot \cdots \odot u'_n) = \\ & \sum \mu^{[l-1]} \epsilon(\sigma) (\exp(f) \circ \exp(g))_{i_1, \dots, i_k}^{t_1, \dots, t_l} (u'_{\sigma(1)} \odot \cdots \odot u'_{\sigma(i_1)} \otimes \cdots \otimes (u'_{\sigma(n-i_k+1)} \odot \cdots \odot u'_{\sigma(n)}), \end{aligned}$$

where

$$(27) \quad (\exp(f) \circ \exp(g))_{i_1, \dots, i_k}^{t_1, \dots, t_l} =: \sum \frac{1}{k! l!} \begin{pmatrix} f_{s_1}^{t_1} \otimes \cdots \otimes f_{s_l}^{t_l} \\ \Psi_{j_1, \dots, j_k}^{s_1, \dots, s_l} \\ g_{i_1}^{j_1} \otimes \cdots \otimes g_{i_k}^{j_k} \end{pmatrix}$$

with the summation as in (25). The crucial difference against (25) is however that (27) involves the entire $\Psi_{j_1, \dots, j_k}^{s_1, \dots, s_l}$ not only its connected part.

Now the theorem follows from the principle standard in the theory of Feynman diagrams^{||} that the logarithm singles out connected components of graphs or, equivalently, that the exponential assembles graphs from their connected components. Let us explain what this principle says in our case. Calculating the exponential of $f \diamond g$ using formula (26) involves, instead of (27), expressions like

$$(28) \quad \sum \frac{1}{u! k_1! l_1! \cdots k_u! l_u!} \begin{pmatrix} f_{s_1^1}^{t_1^1} \otimes \cdots \otimes f_{s_{l_1}^1}^{t_{l_1}^1} \\ {}^1\Psi_{j_1^1, \dots, j_{k_1}^1}^{s_1^1, \dots, s_{l_1}^1} \\ g_{i_1^1}^{j_1^1} \otimes \cdots \otimes g_{i_{k_1}^1}^{j_{k_1}^1} \end{pmatrix} \otimes \cdots \otimes \begin{pmatrix} f_{s_1^u}^{t_1^u} \otimes \cdots \otimes f_{s_{l_u}^u}^{t_{l_u}^u} \\ {}^1\Psi_{j_1^u, \dots, j_{k_u}^u}^{s_1^u, \dots, s_{l_u}^u} \\ g_{i_1^u}^{j_1^u} \otimes \cdots \otimes g_{i_{k_u}^u}^{j_{k_u}^u} \end{pmatrix}$$

with some $u \geq 1$, $l_1 + \cdots + l_u = l$, $k_1 + \cdots + k_u = k$, and

$$t_1^1 + \cdots + t_{l_1}^1 + \cdots + t_1^u + \cdots + t_{l_u}^u = m \quad \text{and} \quad i_1^1 + \cdots + i_{k_1}^1 + \cdots + i_1^u + \cdots + i_{k_u}^u = n.$$

In fact, formula (28) expresses the right hand side of (27) via products of its ‘connected’ components. The seeming discrepancy between the coefficients

$$(29) \quad \frac{1}{k! l!} \quad \text{in (27) and} \quad \frac{1}{u! k_1! l_1! \cdots k_u! l_u!} \quad \text{in (28)}$$

is explained as follows. While in (27) the connected components are entangled, in (28) they are separated. However, thanks to the commutativity of symmetric algebras, the entangled components can be separated using suitable permutations on the output and input sides. It is straightforward though tedious to check that, with the coefficients (29), the corresponding terms appear with the same multiplicity. $\square$

Corollary 33. *IBL $_{\infty}$ -algebras with morphisms (17) form a subcategory IBL $_{\infty}$ of $\mathbf{MV}_{\mathbf{k}[[h]]}$.*

^{||}See e.g. [1, p. 119].

Proof. It is simple to verify that a map as in (17) satisfies (18) if and only if its $\binom{m}{n}$ -component f_n^m is divisible by $\hbar^{n-1}$ for each $m \geq 0$ and $n \geq 1$, i.e. if

$$f_n^m = \hbar^{n-1} \phi_n^m, \quad m \geq 0, \quad n \geq 1,$$

for some $\phi_n^m : S^n(U') \rightarrow S^m(U'')[[h]]$. Let therefore

$$f : S(U'')[[h]] \rightarrow S(U''')[[h]] \quad \text{and} \quad g : S(U')[[h]] \rightarrow S(U'')[[h]]$$

be such maps. We must prove that each $\binom{m}{n}$ -component $(f \diamond g)_n^m$ of their MV-composition (19) with $n \geq 1$ is divisible by $\hbar^{n-1}$. This clearly happens if the expression (25) is, for $n = i_1 + \dots + i_k$, divisible by $\hbar^{n-1}$.

Since each $f_{s_b}^{t_b}$ is divisible by $\hbar^{s_b-1}$ if $s_b \geq 1$, $1 \leq b \leq l$, and each $g_{i_a}^{j_a}$ is divisible by $\hbar^{i_a-1}$ if $i_a \geq 1$, $1 \leq a \leq k$ by assumption, (25) is divisible by

$$\hbar^{i_1 + \dots + i_k + s_1 + \dots + s_l - (\bar{k} + \bar{l})} = \hbar^{n + s_1 + \dots + s_l - (\bar{k} + \bar{l})},$$

where $\bar{k}$ is the number of a 's for which $i_a \neq 0$ (resp. $\bar{l}$ the number of b 's for which $s_b \neq 0$). Thus (25) is divisible by $\hbar^{n-1}$ if

$$(30) \quad n + s_1 + \dots + s_l - (\bar{k} + \bar{l}) \geq n - 1.$$

Since $\bar{k} \leq k$ and $\bar{l} \leq l$, (30) would follow from

$$n + s_1 + \dots + s_l - (k + l) \geq n - 1,$$

which is the same as

$$s_1 + \dots + s_l - (k + l) + 1 \geq 0.$$

Notice that (25) is the sum over unshuffles κ whose associated graphs Γ are connected. The number of vertices V of such a graph clearly equals $k + l$ while its number E of edges is $s_1 + \dots + s_l$. We therefore need to prove that

$$(31) \quad E - V + 1 \geq 0.$$

Since Γ connected, $V - E = 1 - b_1(\Gamma)$ by Euler's theorem or, equivalently, $E - V + 1 = b_1(\Gamma)$, where $b_1(\Gamma)$ is the first Betti number of Γ . As $b_1(\Gamma) \geq 0$, (31) immediately follows. $\square$

6. THE QUANTUM MASTER EQUATION

Definition 34. Let V be an MV-algebra over R with a maximal ideal $\mathfrak{m}$ as in Definition 1. The *quantum master equation (QME)* in V is the equation

$$(32) \quad \Delta e^S = 0,$$

for a degree-0 element $S \in V \widehat{\otimes} \mathfrak{m}$. We denote by $\text{QME}_V(R)$ the set of all solutions of the quantum master equation in V .

In (32), e^S is the exponential in the graded commutative associative algebra $V \widehat{\otimes} R$. The existence of the exponential is guaranteed by the completeness of R .

Example 35 (Münster-Sachs' formulation). Let A be an IBL_∞ -algebra as in Example 10. Münster and Sachs consider $S \in S(U)[[\hbar]]$ of the form

$$(33) \quad S = \sum_{k \geq 2} \hbar^{k-1} c_k$$

with some $c_k \in \bigoplus_{1 \leq n \leq k} S^n(U)$.

Remark 36. Münster and Sachs allowed $k = 1$ in (33), which would be inaccurate, unless there were reasons to guarantee convergence of e^S – see the next example.

Example 37 (Formulation of [2]). Let A be a BV_∞ -algebra as in Example 9. Consider the functional field

$$\mathbb{K} := \mathbb{k}((\hbar)) = \mathbb{k}[[\hbar]][\hbar^{-1}].$$

For an independent symbol λ , $\mathbb{k}[[\hbar]][[\lambda]] = \mathbb{k}[[\hbar, \lambda]]$ is a complete local ring with residue field $\mathbb{k}$ and $\mathbb{K}[[\lambda]] := \mathbb{k}((\hbar))[[\lambda]]$ is a complete local ring with residue field $\mathbb{K}$. Extending Remark 3, we may consider A as an MV-algebra over $\mathbb{k}[[\hbar, \lambda]]$ or over $\mathbb{k}((\hbar))[[\lambda]]$.

The work [2] treated the quantum master equation for S of the form $S = \tilde{S}/\hbar$, with some $\tilde{S} \in \lambda A[[\hbar]][[\lambda]]$ or $\tilde{S} \in \lambda A((\hbar))[[\lambda]]$. Here we work in our more general setup and consider $\tilde{S} \in A((\hbar)) \hat{\otimes} \mathfrak{n}$, where $\mathfrak{n}$ is the maximal ideal in a complete local ring T with residue field $\mathbb{k}$.

Theorem 38 ([3]). *In the situation of Example 37, the equation $\Delta e^{\tilde{S}/\hbar} = 0$ for $\tilde{S} \in A((\hbar)) \hat{\otimes} \mathfrak{n}$ of degree zero is equivalent to*

$$\Delta \tilde{S} + \frac{1}{2!} l_2(\tilde{S}, \tilde{S}) + \frac{1}{3!} l_3(\tilde{S}, \tilde{S}, \tilde{S}) + \cdots = 0,$$

where l_n for each $n \geq 2$ is the higher derived bracket:

$$l_n(a_1, \dots, a_n) := \frac{1}{\hbar^{n-1}} \Phi_n^\Delta(a_1, \dots, a_n) = \sum_{k=1}^{\infty} \hbar^{k-1} \Phi_n^{\Delta_{k+n-1}}(a_1, \dots, a_n),$$

with $\Phi_n^?$ being defined in (3).

Proof. Let Ad denote the adjoint action $\text{Ad}_g Y := gYg^{-1}$ of the group $\text{GL}(A)$ of invertible, degree-zero linear maps $g : A \rightarrow A$ on the Lie algebra $\mathfrak{gl}(A) := \text{Lin}_{\mathbb{k}}(A, A)$ of all degree-zero linear maps $Y : A \rightarrow A$. Let ad be the adjoint action $\text{ad}_X Y := [X, Y]$ of $\mathfrak{gl}(A)$ on itself. Then, for $X \in \mathfrak{gl}(A) \hat{\otimes} \mathfrak{n}$, we have

$$(34) \quad \text{Ad}_{e^X} = e^{\text{ad}_X}.$$

Given a degree-zero element $\tilde{S} \in A((\hbar)) \hat{\otimes} \mathfrak{n}$, apply Equation (34) to the operator $X = L_{(-\tilde{S}/\hbar)}$ of left multiplication by $-\tilde{S}/\hbar$. We get

$$\Delta \circ e^{\tilde{S}/\hbar} = e^{\tilde{S}/\hbar} \sum_{n=0}^{\infty} \frac{\text{ad}_{(-\tilde{S}/\hbar)}^n}{n!} \Delta = e^{\tilde{S}/\hbar} \sum_{n=0}^{\infty} \frac{[[\dots [\Delta, L_{\tilde{S}/\hbar}], \dots], L_{\tilde{S}/\hbar}]}{n!},$$

where in the last formula there are exactly n iterated commutators with $L_{\tilde{S}/\hbar}$. Applying both sides of this equation to $1 \in A$ and using Equation (3), we obtain

$$\Delta(e^{\tilde{S}/\hbar}) = \frac{e^{\tilde{S}/\hbar}}{\hbar} \left(\Delta\tilde{S} + \frac{1}{2!}l_2(\tilde{S}, \tilde{S}) + \frac{1}{3!}l_3(\tilde{S}, \tilde{S}, \tilde{S}) + \cdots \right),$$

whence the result. $\square$

Theorem 39. *There is a natural one-to-one correspondence between solutions S of the QME (32) in V and MV-morphisms $s \in \text{MV}_R(\mathbf{k}, V)$, i.e. one has a natural isomorphism*

$$(35) \quad \text{QME}_V(R) \cong \text{MV}_R(\mathbf{k}, V),$$

given by $s(1) = S$. In (35), the MV structure on $\mathbf{k}$ is defined by $\delta(1) := 1 \otimes 1$ and $\Delta = 0$, as in Example 4.

Remark 40 (Geometric interpretation). Using the geometric interpretation of MV-morphisms in Remark 22, we can interpret the proposition as stating that a solution of the QME is a family X of MV-manifolds over $B = \text{Spec } R$, or rather a family of MV-manifold structures on the trivial fiber bundle $X = \text{Spec } V \hat{\times} B$. One can also view a solution as a B -point $B \rightarrow X^*$ of the dual MV-manifold.

Proof of Theorem 39. By definition, $\text{Lin}_R^0(\mathbf{k}, V)$ consists of R -linear maps

$$(36) \quad s : \mathbf{k} \hat{\otimes} R \cong R \rightarrow V \hat{\otimes} R$$

such that $s(1) \in V \hat{\otimes} \mathbf{m}$. Under the canonical isomorphism

$$\text{Lin}_R(\mathbf{k}, V) \cong V \hat{\otimes} R, \quad s \in \text{Lin}_R(\mathbf{k}, V) \longmapsto s(1) \in V \hat{\otimes} R,$$

maps with this property tautologically correspond to $V \hat{\otimes} \mathbf{m}$.

A map s in (36) is an MV-morphism if and only if $\Delta \circ \exp(s) = 0$. Since $\Delta \circ \exp(s)$ is R -linear, this happens if and only if $\Delta \exp(s)(1) = 0$. As $1 \in \mathbf{k}$ is group-like,

$$\Delta \exp(s)(1) = \Delta e^{s(1)} = \Delta e^S$$

by Example 13. $\square$

Let V be an MV-algebra over $\mathbf{k}[[\hbar]]$ and T a complete local ring with the maximal ideal $\mathbf{n}$ and residue field $\mathbf{k}$. Extend the MV structure on V linearly, as in Remark 3, to an MV structure on the same V over $\mathbf{k}[[\hbar]] \hat{\otimes} T$ and $\mathbf{k}((\hbar)) \hat{\otimes} T$, regarded as complete local rings with residue fields $\mathbf{k}$ and $\mathbf{K} = \mathbf{k}((\hbar))$, respectively.

In the BV formalism of theoretical physics [13] and applications to deformation theory [8, 10, 18], it is important to consider the QME for elements S of the form $S = \tilde{S}/\hbar$ for $\tilde{S} \in V((\hbar)) \hat{\otimes} \mathbf{n}$ or $\tilde{S} \in V[[\hbar]] \hat{\otimes} \mathbf{n}$. We relate these types of solutions to MV-morphisms below.

Corollary 41. (i) *There is a natural one-to-one correspondence between solutions $\tilde{S} \in V((\hbar)) \hat{\otimes} \mathfrak{n}$ of the quantum master equation*

$$(37) \quad \Delta e^{\tilde{S}/\hbar} = 0$$

and MV-morphisms $s \in \mathbf{MV}_{\mathbb{K} \hat{\otimes} T}(\mathbb{K}, V)$, i.e. one has a natural isomorphism

$$(38) \quad \{\tilde{S} \in V((\hbar)) \hat{\otimes} \mathfrak{n} \mid \Delta e^{\tilde{S}/\hbar} = 0\} = \hbar \cdot \text{QME}_{V((\hbar))}(\mathbb{K} \hat{\otimes} T) \xrightarrow{\sim} \mathbf{MV}_{\mathbb{K} \hat{\otimes} T}(\mathbb{K}, V((\hbar))),$$

given by $s(1) = \tilde{S}/\hbar$.

(ii) *There is a natural one-to-one correspondence between solutions $\tilde{S} \in V[[\hbar]] \hat{\otimes} \mathfrak{n}$ of the quantum master equation*

$$(39) \quad \Delta e^{\tilde{S}/\hbar} = 0$$

and a certain subset of MV-morphisms $s \in \mathbf{MV}_{\mathbb{K} \hat{\otimes} T}(\mathbb{K}, V((\hbar)))$, namely one has a natural isomorphism

$$(40) \quad \{\tilde{S} \in V[[\hbar]] \hat{\otimes} \mathfrak{n} \mid \Delta e^{\tilde{S}/\hbar} = 0\} \xrightarrow{\sim} \mathbf{MV}_{\mathbb{K} \hat{\otimes} T}^s(\mathbb{K}, V((\hbar))),$$

given by $s(1) = \tilde{S}/\hbar$, where $\mathbf{MV}_{\mathbb{K} \hat{\otimes} T}^s(\mathbb{K}, V((\hbar)))$ denotes the set of MV-morphisms s whose value $s(1)$ at $1 \in \mathbb{K} \hat{\otimes} T$ has at most a simple pole at $\hbar = 0$.

Remark 42 (Geometric interpretation). Now, as we have two local algebras, geometric interpretation becomes subtler. Let D denote $\text{Spec } k[[\hbar]]$, often called the *formal disk*, $\mathring{D} = \text{Spec } \mathbb{K}$ be the deleted formal disk, and $C = \text{Spec } T$. An MV-algebra over $k[[\hbar]]$ has been interpreted as an MV-manifold over D , which may be restricted to an MV-manifold over $\mathring{D}$. Let us denote it $X_0 \rightarrow \mathring{D}$. Then the isomorphism (38) interprets a solution of the modified QME (37) as a family of MV-manifolds extending the family $X_0 \rightarrow \mathring{D}$ to the base $\mathring{D} \hat{\times} C$:

$$\begin{array}{ccc} X_0 & \longrightarrow & X \\ \downarrow & & \downarrow \\ \mathring{D} & \longrightarrow & \mathring{D} \hat{\times} C. \end{array}$$

In the dual interpretation, the isomorphism (38) asserts that the dual MV-manifold $X_0^* \rightarrow \mathring{D}$ is an MV-manifold representing the functor solutions of QME (37) or the “moduli MV-space” of solutions of the QME. Indeed, a solution of (37) may be interpreted as an MV-morphism

$$\begin{array}{ccc} \mathring{D} \hat{\times} C & \longrightarrow & X_0^* \\ & \searrow & \downarrow \\ & & \mathring{D}. \end{array}$$

Theorem 39 treats solutions of the QME (32) as morphisms in the category of MV-algebras over R . Thus, composition of MV-morphisms can be used to transfer these solutions over MV-morphisms. More specifically, let $f \in \mathbf{MV}_R(V', V'')$ be a morphism and $S \in V' \hat{\otimes} \mathfrak{m}$ a solution of the QME in V' . Theorem 39 translates S into a morphism $s \in \mathbf{MV}_R(k, V')$. Then

$f \diamond s$ will be a morphism in $\text{MV}_R(\mathbf{k}, V'')$ therefore, by Theorem 39 again, $(f \diamond s)(1)$ will solve the QME in V'' . Since $1 \in \mathbf{k}$ is group-like, one moreover has

$$(f \diamond s)(1) = \log((\exp(f) \circ \exp(s))(1)) = \log(\exp(f)(e^{s(1)})) = \log(\exp(f)(e^S)),$$

the ‘actual’ logarithm of $\exp(f)(e^S) \in V'' \widehat{\otimes} \mathbf{m}$.

Definition 43. The element

$$f_!(S) := \log(\exp(f)(e^S)) \in \text{QME}_{V''}(R)$$

is the *push-forward* of the solution $S \in \text{QME}_{V'}(R)$.

Theorem 44 (Cf. [2] for the case $V' = S(U')$). *Assume that V' is a bialgebra, i.e. the standard compatibility between the multiplication and the comultiplication is fulfilled. Suppose moreover that $S \in P(V') \widehat{\otimes} \mathbf{m}$, where $P(V')$ is the subspace of primitive elements in V' . Then*

$$f_!(S) = f(e^S).$$

Proof. Observe that e^S is a group-like element in $V' \widehat{\otimes} R$. Indeed,

$$\delta'(e^S) = e^{\delta'(S)} = e^{(S \otimes_R 1 + 1 \otimes_R S)} = e^{S \otimes_R 1} e^{1 \otimes_R S} = e^S \otimes_R e^S,$$

where we used the fact that δ' is an algebra morphism and that $S \otimes_R 1$ commutes with $1 \otimes_R S$ in $(V' \widehat{\otimes} R) \otimes_R (V' \widehat{\otimes} R)$. The rest follows from the observations in Example 13. $\square$

REFERENCES

- [1] D. Bessis, C. Itzykson, and J. B. Zuber. Quantum field theory techniques in graphical enumeration. *Adv. in Appl. Math.*, 1(2):109–157, 1980.
- [2] D. Bashkirov and A. A. Voronov. The BV formalism for L_∞ -algebras. Preprint IHES/M/14/36, [arXiv:1410.6432 \[math.QA\]](#).
- [3] Ch. Braun and A. Lazarev. Homotopy BV algebras in Poisson geometry. *Trans. Moscow Math. Soc.*, pages 217–227, 2013.
- [4] R. Campos, S. Merkulov, T. Willwacher. The Frobenius properad is Koszul. Preprint [arXiv:1402.4048 \[math.QA\]](#).
- [5] K. Cieliebak, K. Fukaya, and J. Latschev. Homological algebra related to surfaces with boundaries. Preprint [arXiv:1508.02741 \[math.QA\]](#).
- [6] K. Cieliebak and J. Latschev. The role of string topology in symplectic field theory. In *New perspectives and challenges in symplectic field theory*, volume 49 of *CRM Proc. Lecture Notes*, pages 113–146. Amer. Math. Soc., Providence, RI, 2009.
- [7] G. C. Drummond-Cole, J. Terilla, and T. Tradler. Algebras over $\Omega(\text{coFrob})$. *J. Homotopy Relat. Struct.*, 5(1):15–36, 2010.
- [8] D. Iacono. Deformations and obstructions of pairs (X, D) . Preprint [arXiv:1302.1149 \[math.AG\]](#).
- [9] Ch. Kassel. *Quantum groups*, volume 155 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 1995.
- [10] L. Katzarkov, M. Kontsevich, and T. Pantev. Hodge theoretic aspects of mirror symmetry. In *From Hodge theory to integrability and TQFT tt*-geometry*, volume 78 of *Proc. Sympos. Pure Math.*, pages 87–174. Amer. Math. Soc., Providence, RI, 2008.
- [11] O. Kravchenko. Deformations of Batalin-Vilkovisky algebras. In *Poisson geometry (Warsaw, 1998)*, volume 51 of *Banach Center Publ.*, pages 131–139. Polish Acad. Sci., Warsaw, 2000.
- [12] T. Lada and M. Markl. Strongly homotopy Lie algebras. *Comm. Algebra*, 23(6):2147–2161, 1995.

- [13] A. Losev. From Berezin integral to Batalin-Vilkovisky formalism: a mathematical physicist's point of view. In *Felix Berezin. Life and death of the mastermind of supermathematics*, pages 3–30. Edited by M. Shifman. World Scientific Publishing Co. Pte. Ltd., Hackensack, NJ, 2007.
- [14] M. Markl. *Deformation theory of algebras and their diagrams*, volume 116 of *CBMS Regional Conference Series in Mathematics*. Published for the Conference Board of the Mathematical Sciences, Washington, DC, 2012.
- [15] M. Markl. On the origin of higher braces and higher-order derivations. *J. Homotopy Relat. Struct.*, (10)(3):637–667, 2015.
- [16] K. Münster and I. Sachs. Quantum open-closed homotopy algebra and string field theory. *Comm. Math. Phys.*, 321(3):769–801, 2013.
- [17] J. Terilla. Quantizing deformation theory. In *Deformation spaces*, volume E40 of *Aspects Math.*, pages 135–141. Vieweg + Teubner, Wiesbaden, 2010.
- [18] J. Terilla. Smoothness theorem for differential BV algebras. *J. Topol.*, 1(3):693–702, 2008.

MATHEMATICAL INSTITUTE OF THE ACADEMY, ŽITNÁ 25, 115 67 PRAGUE 1, THE CZECH REPUBLIC

FACULTY OF MATHEMATICS AND PHYSICS, CHARLES UNIVERSITY, 186 75 SOKOLOVSKÁ 83, PRAGUE 8, THE CZECH REPUBLIC

E-mail address: markl@math.cas.cz

SCHOOL OF MATHEMATICS, UNIVERSITY OF MINNESOTA, 206 CHURCH ST. S.E., MINNEAPOLIS, MN 55455, USA

KAVLI IPMU (WPI), UTIAS, THE UNIVERSITY OF TOKYO, KASHIWA, CHIBA 277-8583, JAPAN

E-mail address: voronov@math.umn.edu